

ORT1273

DNA ENCODING HUMAN SERINE PROTEASE D-G

5 BACKGROUND OF THE INVENTION

Members of the trypsin/chymotrypsin-like (S1) serine protease family play pivotal roles in a multitude of diverse physiological processes, including digestive processes and regulatory amplification cascades through the proteolytic activation of inactive zymogen precursors. In many instances protease substrates within these cascades are themselves the inactive form, or zymogen, of a "downstream" serine protease. Well-known examples of serine protease-mediated regulation include blood coagulation, (Davie, et al. (1991). *Biochemistry* 30:10363-70), kinin formation (Proud and Kaplan (1988). *Ann Rev Immunol* 6: 49-83) and the complement system (Reid and Porter (1981). *Ann Rev Biochemistry* 50:433-464). Although these proteolytic pathways have been known for sometime, it is likely that the discovery of novel serine protease genes and their products will enhance our understanding of regulation within these existing cascades, and lead to the elucidation of entirely novel protease networks.

Proteases are used in non-natural environments for various commercial purposes including laundry detergents, food processing, fabric processing and skin care products. In laundry detergents, the protease is employed to break down organic, poorly soluble compounds to more soluble forms that can be more easily dissolved in detergent and water. In this capacity the protease acts as a "stain remover." Examples of food processing include tenderizing meats and producing cheese.

Proteases are used in fabric processing, for example, to treat wool in order prevent fabric shrinkage. Proteases may be included in skin care products to remove scales on the skin surface that build up due to an imbalance in the rate of desquamation. Common proteases used in some of these applications are derived from prokaryotic or eukaryotic cells that are easily grown for industrial manufacture of their enzymes, for example a common species used is *Bacillis* as described in United States patent

5,217,878. Alternatively, United States Patent 5,278,062 describes serine proteases isolated from a fungus, *Tritirachium album*, for use in laundry detergent compositions. Unfortunately use of some proteases is limited by their potential to

5 cause allergic reactions in sensitive individuals or by reduced efficiency when used in a non-natural environment. It is anticipated that protease proteins derived from non-human sources would be more likely to induce an immune response in a sensitive individual. Because of these limitations, there is a need for alternative proteases that are less immunogenic to sensitive individuals and/or provides efficient proteolytic activity in a non-natural environment. The advent of recombinant technology allows expression of any species' proteins in a host suitable for industrial manufacture.

Herein we describe a novel serine protease isolated from small intestine termed D-G. The deduced amino acid sequence encodes a polypeptide of 435 amino acids. Interestingly, the sequence contains a hydrophobic stretch of amino acids which is a putative transmembrane near the NH₂-terminus. Thus, this serine protease is thought to be synthesized as a type II integral membrane protein. Alignment with other well characterized serine proteases clearly indicates that it is a member of the S1 serine protease family with the catalytic triad residing within the C-terminal half of the molecule. The protease D-G deduced amino acid sequence is most similar to the cloned serine proteases TMPRSS2 (Paoloni-Giacobino et al. (1997). *Genomics* 44:309-320) and hepsin (Leytus et al. (1988). *Biochemistry* 27:1067-74), which are also type II integral membrane proteases. We have found that the protease D-G mRNA is widely expressed in several tissues throughout the body including epidermis, fibroblasts, keratinocytes, colon, small intestine, stomach, lung, kidney, bone marrow, lymph node, thymus, ovary, prostate, uterus and spinal cord. Altered expression or regulation of this enzyme may be responsible for any one of a number of pathological conditions in these tissues. Furthermore, an up-regulation whereby under normal physiological conditions protease D-G mRNA is not expressed, and therefore undetected, but in the pathogenic condition it is markedly elevated could potentially result in initiating or exacerbation of certain diseased states. We

ORT1273

expressed a soluble form of this novel human protease by inserting the portion of the protease D-G cDNA, encoding the catalytic domain, in a zymogen activation construct designed to permit the generic activation of heterologous serine protease catalytic domains. The result is an active preparation of protease D-G that has an activity against a subset of amidolytic substrates. Isolation of purified, enzymatically active protease D-G allows the protein to be used directly, for example to discover chemical modulators of the enzyme or as an additive in commercial products. Because protease D-G is derived from a human host, it is less likely to induce an allergic reaction in sensitive individuals, and therefore protease D-G may also be useful for formulation of compositions for laundry detergents and skin care products.

SUMMARY OF THE INVENTION

A DNA molecule encoding protease D-G has been cloned and characterized and it represents a novel serine protease. Using a recombinant expression system functional DNA molecules encoding the protease have been isolated. The biological and structural properties of these proteins are disclosed, as is the amino acid and nucleotide sequence. The recombinant DNA molecules, and portions thereof, are useful for isolating homologues of the DNA molecules, identifying and isolating genomic equivalents of the DNA molecules, and identifying, detecting or isolating mutant forms of the DNA molecules. The recombinant protein is useful to identify modulators of functional protease D-G. Modulators identified in the assays disclosed herein may be useful as therapeutic agents for cancer, skin disorders, neuropathic pain, inflammatory, or coagulation diathesis/thrombosis.

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 A - The nucleotide (SEQ.ID.NO.:1) of the novel protease D-G cDNA is shown.

Figure 1 B - The amino acid sequence (SEQ.ID.NO.:2) of the novel protease D-G cDNA is shown.

5

Figure 2 - The phylogenetic tree of the protease D-G amino acid sequence relative to other S1 serine proteases is shown.

Figure 3 - PCR-based tissue distribution indicates that the protease D-G mRNA is restricted. Autoradiograms of gels are shown with the position of the D-G specific PCR product, as detected by the hybridization of a labeled nested probe, which was resolved following electrophoresis from the free probe (F.P.). The cDNA libraries of tissues and cell lines analyzed are as indicated.

Figure 4 A & B- The nucleotide (SEQ.ID.NO.:8) and amino acid (SEQ.ID.NO.:9) sequences of the protease D-G catalytic domain in the zymogen activation construct are shown.

Figure 5 - Polyacrylamide gel and Western blot analyses of the purified recombinant protease PFEK-protease D-G-6XHIS. Shown is the polyacrylamide gel containing samples of the novel serine protease PFEK-protease D-G-6XHIS stained with Coomassie Brilliant Blue (lanes 2 and 3). The relative molecular masses are indicated by the positions of protein standards (lane 1). In the indicated lanes, the purified zymogen was either untreated (-) or digested (+) with enterokinase (EK) which was used to cleave and activate the zymogen of lane 1 into its active form of increased mobility shown in lane 2. Lanes 4 and 5 indicate the Western blot of the corresponding gel lanes 1 and 2, probed with the anti-FLAG MoAb M2. This demonstrates the quantitative cleavage of the expressed and purified zymogen to generate the processed and activated protease. Since the FLAG epitope is located just upstream of the of the EK pro sequence, cleavage with EK generates a FLAG-containing polypeptide which is

30

Lanes 4 and 5 indicate the Western blot of the corresponding gel lanes 1 and 2, probed with the anti-FLAG MoAb M2. This demonstrates the quantitative cleavage of the expressed and purified zymogen to generate the processed and activated protease. Since the FLAG epitope is located just upstream of the of the EK pro sequence, cleavage with EK generates a FLAG-containing polypeptide which is

ORT1273

too small to be retained in the polyacrylamide gel, and is therefore not detected in the +EK lane.

Figure 6 - Functional amidolytic activities of the recombinant protease D-G-6XHIS expressed, purified and activated from the activation construct were determined using the indicated chromogenic substrates.

DETAILED DESCRIPTION

Definitions

10 The term "protein domain" as used herein refers to a region of a protein that may have a particular three-dimensional structure which may be independent from the remainder of the protein. This structure may maintain a particular activity associated with the domain's function within the protein including enzymatic activity, creation of a recognition motif for another molecule, or provide necessary structural

15 components for a protein to exist in a particular environment. Protein domains are usually evolutionarily conserved regions of proteins, both within a protein family and within protein superfamilies that perform similar functions. The term "protein superfamily" as used herein refers to proteins whose evolutionary relationship may not be entirely established or may be distant by accepted phylogenetic standards, but

20 show similar three dimensional structure or display unique consensus of critical amino acids. The term "protein family" as used herein refers to proteins whose evolutionary relationship has been established by accepted phylogenic standards.

The term "fusion protein" as used herein refers to protein constructs that are the result of combining multiple protein domains or linker regions for the purpose of

25 gaining the combined functions of the domains or linker regions. This is may be accomplished by molecular cloning of the nucleotide sequences encoding such domains to produce a new polynucleotide sequence that encodes the desired fusion protein. Alternatively, creation of a fusion protein may be accomplished by chemically joining two proteins.

30 The term "linker region" or "linker domain" or similar such descriptive terms as used herein refers to polynucleotide or polypeptide sequence that are used in the

ORT1273

construction of a cloning vector or fusion protein. Functions of a linker region can include introduction of cloning sites into the nucleotide sequence, introduction of a flexible component or space-creating region between two protein domains, or creation of an affinity tag for specific molecule interaction. A linker region may be introduced into a fusion protein resulting from choices made during polypeptide or nucleotide sequence construction.

The term "cloning site" or "polycloning site" as used herein refers to a region of the nucleotide sequence that has one or more available restriction endonuclease consensus cleavage sequences. These nucleotide sequences may be used for a variety of purposes, including but not limited to introduction into DNA vectors to create novel fusion proteins, or to introduce specific site-directed mutations. It is well known by those of ordinary skill in the art that cloning sites can be engineered at a desired location by silent mutations, conserved mutation, or introduction of a linker region that contains desired restriction enzyme consensus sequences. It is also well known by those of ordinary skill in the art that the precise location of a cloning site can be engineered into any location in a nucleotide sequence.

The term "tag" as used herein refers to an amino acid sequence or a nucleotide sequence that encodes an amino acid sequence, that facilitates isolation, purification or detection of a protein containing the tag. A wide variety of such tags are known to those skilled in the art, and are suitable for use in the present invention. Suitable tags include, but are not limited to, HA peptide, polyhistidine peptides, biotin / avidin, and other antibody epitope binding sites.

Isolation of protease D-G nucleic acid

25

The present invention relates to DNA encoding the human serine protease D-G which was isolated from cells of small intestine. Protease D-G, as used herein, refers to protein which can specifically function as a protease.

The complete amino acid sequence of protease D-G was not previously known, nor was the complete nucleotide sequence encoding protease D-G known. It is predicted that a wide variety of cells and cell types will contain the described

protease D-G mRNA. Tissues capable of producing protease D-G include, but are not limited to epidermis, fibroblasts, keratinocytes, colon, small intestine, stomach, lung, kidney, bone marrow, lymph node, thymus, ovary, prostate, uterus and spinal cord as we have determined by a sensitive polymerase chain reaction (PCR)-mediated mRNA detection methodology.

Any of a variety of procedures known in the art may be used to molecularly clone protease D-G DNA. These methods include, but are not limited to, direct functional expression of the protease D-G genes following the construction of a protease D-G-containing cDNA library in an appropriate expression vector system. Another method is to screen protease D-G-containing cDNA library constructed in a bacteriophage or plasmid shuttle vector with a labelled oligonucleotide probe designed from the amino acid sequence of the protease D-G subunits. An additional method consists of screening a protease D-G-containing cDNA library constructed in a bacteriophage or plasmid shuttle vector with a partial cDNA encoding the protease D-G protein. This partial cDNA is obtained by the specific PCR amplification of protease D-G DNA fragments through the design of degenerate oligonucleotide primers from the amino acid sequence of the purified protease D-G protein.

Another method is to isolate RNA from protease D-G-producing cells and
25 translate the RNA into protein via an *in vitro* or an *in vivo* translation system. The
translation of the RNA into a peptide a protein will result in the production of at
least a portion of the protease D-G protein which can be identified by, for
example, immunological reactivity with an anti-protease D-G antibody or by
biological activity of protease D-G protein. In this method, pools of RNA isolated
30 from protease D-G-producing cells can be analyzed for the presence of an RNA

ORT1273

that encodes at least a portion of the protease D-G protein. Further fractionation of the RNA pool can be done to purify the protease D-G RNA from non-protease D-G RNA. The peptide or protein produced by this method may be analyzed to provide amino acid sequences which in turn are used to provide primers for production of protease D-G cDNA, or the RNA used for translation can be analyzed to provide nucleotide sequences encoding protease D-G and produce probes for this production of protease D-G cDNA. This method is known in the art and can be found in, for example, Maniatis, T., Fritsch, E.F., Sambrook, J. in Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 1989.

It is readily apparent to those skilled in the art that other types of libraries, as well as libraries constructed from other cells or cell types, may be useful for isolating protease D-G-encoding DNA. Other types of libraries include, but are not limited to, cDNA libraries derived from other cells, from organisms other than protease D-G, and genomic DNA libraries that include YAC (yeast artificial chromosome) and cosmid libraries.

It is readily apparent to those skilled in the art that suitable cDNA libraries may be prepared from cells or cell lines which have protease D-G activity. The selection of cells or cell lines for use in preparing a cDNA library to isolate protease D-G cDNA may be done by first measuring cell associated protease D-G activity using the measurement of protease D-G-associated biological activity or a ligand binding assay.

Preparation of cDNA libraries can be performed by standard techniques well known in the art. Well known cDNA library construction techniques can be found for example, in Maniatis, T., Fritsch, E.F., Sambrook, J., Molecular Cloning: A Laboratory Manual, Second Edition (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989).

It is also readily apparent to those skilled in the art that DNA encoding protease D-G may also be isolated from a suitable genomic DNA library.

ORT1273

Construction of genomic DNA libraries can be performed by standard techniques well known in the art. Well known genomic DNA library construction techniques can be found in Maniatis, T., Fritsch, E.F., Sambrook, J. in *Molecular Cloning: A Laboratory Manual*, Second Edition (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989).

In order to clone the protease D-G gene by the above methods, the amino acid sequence of protease D-G may be necessary. To accomplish this, protease D-G protein may be purified and partial amino acid sequence determined by automated sequenators. It is not necessary to determine the entire amino acid sequence, but the linear sequence of two regions of 6 to 8 amino acids from the protein is determined for the production of primers for PCR amplification of a partial protease D-G DNA fragment.

Once suitable amino acid sequences have been identified, the DNA sequences capable of encoding them are synthesized. Because the genetic code is degenerate, more than one codon may be used to encode a particular amino acid, and therefore, the amino acid sequence can be encoded by any of a set of similar DNA oligonucleotides. Only one member of the set will be identical to the protease D-G sequence but will be capable of hybridizing to protease D-G DNA even in the presence of DNA oligonucleotides with mismatches. The mismatched DNA oligonucleotides may still sufficiently hybridize to the protease D-G DNA to permit identification and isolation of protease D-G encoding DNA. DNA isolated by these methods can be used to screen DNA libraries from a variety of cell types, from invertebrate and vertebrate sources, and to isolate homologous genes.

Purified biologically active protease D-G may have several different physical forms. protease D-G may exist as a full-length nascent or unprocessed polypeptide, or as partially processed polypeptides or combinations of processed polypeptides. The full-length nascent protease D-G polypeptide may be post-translationally modified by specific proteolytic cleavage events that results in the formation of fragments of the full length nascent polypeptide. A fragment, or physical association

ORT1273

of fragments may have the full biological activity associated with protease D-G however, the degree of protease D-G activity may vary between individual protease D-G fragments and physically associated protease D-G polypeptide fragments.

Because the genetic code is degenerate, more than one codon may be used to encode a particular amino acid, and therefore, the amino acid sequence can be encoded by any of a set of similar DNA oligonucleotides. Only one member of the set will be identical to the protease D-G sequence but will be capable of hybridizing to protease D-G DNA even in the presence of DNA oligonucleotides with mismatches under appropriate conditions. Under alternate conditions, the mismatched DNA oligonucleotides may still hybridize to the protease D-G DNA to permit identification and isolation of protease D-G encoding DNA.

DNA encoding protease D-G from a particular organism may be used to isolate and purify homologues of protease D-G from other organisms. To accomplish this, the first protease D-G DNA may be mixed with a sample containing DNA encoding homologues of protease D-G under appropriate hybridization conditions. The hybridized DNA complex may be isolated and the DNA encoding the homologous DNA may be purified therefrom.

Functional derivatives / Variants

It is known that there is a substantial amount of redundancy in the various codons that code for specific amino acids. Therefore, this invention is also directed to those DNA sequences that contain alternative codons that code for the eventual translation of the identical amino acid. For purposes of this specification, a sequence bearing one or more replaced codons will be defined as a degenerate variation. Also included within the scope of this invention are mutations either in the DNA sequence or the translated protein, which do not substantially alter the ultimate physical properties of the expressed protein. For example, substitution of aliphatic amino acids alanine, valine, leucine and isoleucine; interchange of the hydroxyl residues serine and threonine, exchange of the acidic residues aspartic acid and glutamic acid, substitution between the amide residues asparagine and glutamine, exchange of the

basic residues lysine and arginine and variants among the aromatic residues phenylalanine, tyrosine may not cause a change in functionality of the polypeptide. Such substitutions are well known and are described, for instance in Molecular Biology of the Gene, 4th Ed. Benjamin Cummings Pub. Co. by Watson *et al*.

Identity or similarity, as known in the art, are relationships between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. Both identity and similarity can be readily calculated (Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and

Identity or similarity, as known in the art, are relationships between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. Both identity and similarity can be readily calculated (Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and

Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, M. J., Academic Press, 1989.

- 5 J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity and similarity between two polynucleotide or two polypeptide sequences, both terms are well known to skilled artisans (Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and
10 Carillo, H., and Lipman, D., (1988) SIAM J. Applied Math., 48, 1073. Methods commonly employed to determine identity or similarity between sequences include, but are not limited to those disclosed in Carillo, H., and Lipman, D., (1988) SIAM J. Applied Math., 48, 1073. Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity
15 and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCG program package (Devereux, J., et al., (1984) Nucleic Acids Research 12(1), 387), BLASTP, BLASTN, and FASTA (Atschul, S. F. et al., (1990) J. Molec. Biol. 215, 403).
- 20 Polynucleotide(s) generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. Thus, for instance, polynucleotides as used herein refers to, among others, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple- stranded regions, single- and double-
25 stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded, or a mixture of single- and double-stranded regions. In addition, polynucleotide as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions
30 may be from the same molecule or from different molecules. The regions may include

all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term polynucleotide includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with

The term polypeptides, as used herein, refers to the basic chemical structure of polypeptides that is well known and has been described in textbooks and other publications in the art. In this context, the term is used herein to refer to any peptide or protein comprising two or more amino acids joined to each other in a linear chain by peptide bonds. As used herein, the term refers to both short chains, which also commonly are referred to in the art as peptides, oligopeptides and oligomers, for example, and to longer chains, which generally are referred to in the art as proteins, of which there are many types. It will be appreciated that polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids, and that many amino acids, including the terminal amino acids, may be modified in a given polypeptide, either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques which are well known to the art. Even the common modifications that occur naturally in polypeptides are too numerous to list exhaustively here, but they are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known

to those of skill in the art. Among the known modifications which may be present in polypeptides of the present are, to name an illustrative few, acetylation, acylation, ADP- ribosylation, amidation, covalent attachment of flavin, covalent attachment of heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent

5 attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, 10 phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. Such modifications are well known to those of skill and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid 15 residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as, for instance *PROTEINS-- STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as, for example, those provided by Wold, F., *Posttranslational Protein Modifications: Perspectives and 20 Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., (1990) *Meth. Enzymol.* 182, 626-646 and Rattan et al., "Protein Synthesis: Posttranslational Modifications and Aging", (1992) *Ann. N.Y. Acad. Sci.* 663, 48-62. It will be appreciated, as is well known and as noted above, that polypeptides are not 25 always entirely linear. For instance, polypeptides may be generally as a result of posttranslational events, including natural processing event and events brought about by human manipulation which do not occur naturally. Circular, branched and branched circular polypeptides may be synthesized by non-translation natural process and by entirely synthetic methods, as well. Modifications can occur anywhere in a 30 polypeptide, including the peptide backbone, the amino acid side-chains and the

ORT1273

amino or carboxyl termini. In fact, blockage of the amino or carboxyl group in a polypeptide, or both, by a covalent modification, is common in naturally occurring and synthetic polypeptides and such modifications may be present in polypeptides of the present invention, as well. For instance, the amino terminal residue of

5 polypeptides made in *E. coli* or other cells, prior to proteolytic processing, almost invariably will be N-formylmethionine. During post-translational modification of the peptide, a methionine residue at the NH₂-terminus may be deleted. Accordingly, this invention contemplates the use of both the methionine-containing and the methionine-less amino terminal variants of the protein of the invention. The modifications that

10 occur in a polypeptide often will be a function of how it is made. For polypeptides made by expressing a cloned gene in a host, for instance, the nature and extent of the modifications in large part will be determined by the host cell posttranslational modification capacity and the modification signals present in the polypeptide amino acid sequence. For instance, as is well known, glycosylation often does not occur in

15 bacterial hosts such as, for example, *E. coli*. Accordingly, when glycosylation is desired, a polypeptide should be expressed in a glycosylating host, generally a eukaryotic cell. Insect cell often carry out the same posttranslational glycosylations as mammalian cells and, for this reason, insect cell expression systems have been developed to express efficiently mammalian proteins having native patterns of

20 glycosylation, inter alia. Similar considerations apply to other modifications. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. In general, as used herein, the term polypeptide encompasses all such modifications, particularly those that are present in polypeptides

25 synthesized recombinantly by expressing a polynucleotide in a host cell.

Variant(s) of polynucleotides or polypeptides, as the term is used herein, are polynucleotides or polypeptides that differ from a reference polynucleotide or polypeptide, respectively. A variant of the polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant

30 that is not known to occur naturally. (1) A polynucleotide that differs in nucleotide

09607745:063000

sequence from another, reference polynucleotide. Generally, differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical. As noted below, changes in the nucleotide sequence of the variant may be silent. That is, they may not alter the amino acids

polynucleotide. Such nucleotide changes may result in amino acid substitutions,
10 additions, deletions, fusions and truncations in the polypeptide encoded by the

sequence by one or more substitutions, additions, deletions, fusions and truncations, which may be present in any combination. As used herein, a "functional derivative" of protease D-G is a compound that possesses a biological activity (either functional or structural) that is substantially similar to the biological activity of protease D-G.

within the polypeptide with a secondary chemical moiety. Well known cross-linking reagents are useful to react to amino, carboxyl, or aldehyde residues to introduce, for

30 Therefore, if the two molecules possess substantially similar activity, they are

considered to be variants even if the structure of one of the molecules is not found in the other or even if the two amino acid sequences are not identical. The term "analog" refers to a molecule substantially similar in function to either the entire protease D-G molecule or to a fragment thereof. Particularly preferred in this regard

deletions, which do not alter the properties and activities of the gene of SEQ ID NO.:1. Also especially preferred in this regard are conservative substitutions. Most highly preferred are polynucleotides encoding polypeptides having the amino acid sequence of SEQ ID NO.:2, without substitutions.

Further preferred embodiments of the invention are polynucleotides that are at least 70% identical over their entire length to a polynucleotide encoding the polypeptide having the amino acid sequence set out in SEQ ID NO.:2, and polynucleotides which are complementary to such polynucleotides. Alternatively, highly preferred are polynucleotides that comprise a region that is at least 80% identical, more highly preferred are polynucleotides that comprise a region that is at least 90% identical, and among these preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% identity are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the most preferred. The polynucleotides which hybridize to the polynucleotides described herein in a preferred embodiment encode polypeptides which retain substantially the same biological function or activity as the polypeptide characterized by the deduced amino acid sequence of SEQ ID NO.:2. Preferred embodiments in this respect, moreover, are polynucleotides that encode polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by the DNA of SEQ ID NO.:1. The present invention further relates to

ORT1273

polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides that hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences.

As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding the sequences of SEQ ID NO.:1 and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to SEQ ID NO.:1. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less. For example, the coding region of the gene of the invention may be isolated by screening using the known DNA sequence to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the present invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine to which members of the library the probe hybridizes.

The polypeptides of the present invention include the polypeptide of SEQ ID NO.:2 (in particular the mature polypeptide) as well as polypeptides which have at least 70% identity to the polypeptide of SEQ ID NO.:2, preferably at least 80% identity to the polypeptide of SEQ ID NO.:2, and more preferably at least 90% similarity (more preferably at least 90% identity) to the polypeptide of SEQ ID NO.:2 and still more preferably at least 95% similarity (still more preferably at least 97% identity) to the polypeptide of SEQ ID NO.:2 and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids. Representative examples of polypeptide fragments of the invention, include, for example, truncation polypeptides of SEQ ID NO.:2. Truncation polypeptides include polypeptides having the amino acid sequence of SEQ ID NO.:2, or of variants or derivatives thereof, except for

ORT1273

deletion of a continuous series of residues (that is, a continuous region, part or portion) that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or, as in double truncation mutants, deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Also preferred in this aspect of the invention are fragments characterized by structural or functional attributes of the polypeptide characterized by the sequences of SEQ ID NO.:2. Preferred embodiments of the invention in this regard include fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, high antigenic index regions of the polypeptide of the invention, and combinations of such fragments. Preferred regions are those that mediate activities of the polypeptides of the invention. Most highly preferred in this regard are fragments that have a chemical, biological or other activity of the response regulator polypeptide of the invention, including those with a similar activity or an improved activity, or with a decreased undesirable activity.

Recombinant expression of protease D-G

The cloned protease D-G DNA obtained through the methods described herein may be recombinantly expressed by molecular cloning into an expression vector containing a suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to produce recombinant protease D-G protein. Techniques for such manipulations are fully described in Maniatis, T, *et al.*, *supra*, and are well known in the art.

Expression vectors are defined herein as DNA sequences that are required for the transcription of cloned copies of genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express eukaryotic genes in a variety of hosts such as bacteria including *E. coli*, bluegreen algae, plant cells, insect cells, fungal cells including yeast cells, and animal cells.

Specifically designed vectors allow the shuttling of DNA between hosts such as bacteria-yeast or bacteria-animal cells or bacteria-fungal cells or bacteria-invertebrate cells. An appropriately constructed expression vector should contain: an origin of replication for autonomous replication in host cells, selectable

markers, a limited number of useful restriction enzyme sites, a potential for high copy number, and active promoters. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and initiate RNA synthesis. A strong promoter is one that causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses.

A variety of mammalian expression vectors may be used to express recombinant protease D-G in mammalian cells. Commercially available mammalian expression vectors which may be suitable for recombinant protease D-G expression, include but are not limited to, pMAMneo (Clontech), pcDNA3 (Invitrogen), pMC1neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), EBO-pSV2-neo (ATCC 37593) pBPV-1(8-2) (ATCC 37110), pdBPV-MMTneo(342-12) (ATCC 37224), pRSVgpt (ATCC 37199), pRSVneo (ATCC 37198), pSV2-dhfr (ATCC 37146), pUCTag (ATCC 37460), and IZD35 (ATCC 37565).

A variety of bacterial expression vectors may be used to express recombinant protease D-G in bacterial cells. Commercially available bacterial expression vectors which may be suitable for recombinant protease D-G expression include, but are not limited to pET vectors (Novagen) and pQE vectors (Qiagen).

A variety of fungal cell expression vectors may be used to express recombinant protease D-G in fungal cells such as yeast. Commercially available fungal cell expression vectors which may be suitable for recombinant protease D-G expression include but are not limited to pYES2 (Invitrogen) and Pichia expression vector (Invitrogen).

ORT1273

A variety of insect cell expression vectors may be used to express recombinant protease D-G in insect cells. Commercially available insect cell expression vectors which may be suitable for recombinant expression of protease D-G include but are not limited to pBlueBacII (Invitrogen).

- 5 DNA encoding protease D-G may be cloned into an expression vector for expression in a recombinant host cell. Recombinant host cells may be prokaryotic or eukaryotic, including but not limited to bacteria such as E. coli, fungal cells such as yeast, mammalian cells including but not limited to cell lines of human, bovine, porcine, monkey and rodent origin, and insect cells including but not
10 limited to drosophila and silkworm derived cell lines.

- Cell lines derived from mammalian species which may be suitable and which are commercially available, include but are not limited to, CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa
15 (ATCC CCL 2), C127I (ATCC CRL 1616), BS-C-1 (ATCC CCL 26), MRC-5 (ATCC CCL 171), L-cells, and HEK-293 (ATCC CRL1573).

- The expression vector may be introduced into host cells via any one of a number of techniques including but not limited to transformation, transfection, protoplast fusion, lipofection, and electroporation. The expression vector-
20 containing cells are clonally propagated and individually analyzed to determine whether they produce protease D-G protein. Identification of protease D-G expressing host cell clones may be done by several means, including but not limited to immunological reactivity with anti-protease D-G antibodies, and the presence of host cell-associated protease D-G activity.

- 25 Expression of protease D-G DNA may also be performed using *in vitro* produced synthetic mRNA. Synthetic mRNA or mRNA isolated from protease D-G producing cells can be efficiently translated in various cell-free systems, including but not limited to wheat germ extracts and reticulocyte extracts, as well as efficiently translated in cell based systems, including but not limited to

microinjection into frog oocytes, with microinjection into frog oocytes being generally preferred.

20 *Assay methods for protease D-G*

Host cell transfectants and microinjected oocytes may be used to assay both the levels of functional protease D-G activity and levels of total protease D-G protein by the following methods. In the case of recombinant host cells, this involves the co-transfection of one or possibly two or more plasmids, containing the protease D-G DNA encoding one or more fragments encoding the catalytic domain. In the case of oocytes, this involves the co-injection of synthetic RNAs for protease D-G protein. Following an appropriate period of time to allow for expression, cellular protein is metabolically labelled with, for example ³⁵S-methionine for 24 hours, after which cell lysates and cell culture supernatants are

ORT1273

harvested and subjected to immunoprecipitation with polyclonal antibodies directed against the protease D-G protein.

Levels of protease D-G protein in host cells are quantitated by immunoaffinity and/or proteolytic/amidolytic assay techniques. Cells expressing protease D-G can be assayed for the number of protease D-G molecules expressed by measuring the amount of proteolytic/amidolytic activity. Protease D-G-specific affinity beads or protease D-G-specific antibodies are used to isolate for example ³⁵S-methionine labelled or unlabelled protease D-G protein. Labelled protease D-G protein is analyzed by SDS-PAGE. Unlabelled protease D-G protein is detected by Western blotting, ELISA or RIA assays employing protease D-G specific antibodies.

Cell based assays

The present invention provides a whole cell method to detect compound modulation of protease D-G. The method comprises the steps;

- 1) contacting a compound, and a cell that contains functional protease D-G or purifying functional protease D-G, and
- 2) measuring a change in the cell in response or protease D-G activity by the compound.

The amount of time necessary for protease D-G interaction with the compound is empirically determined, for example, by running a time course with a known protease D-G modulator and measuring cellular/activity changes as a function of time.

The term "cell" refers to at least one cell, but includes a plurality of cells appropriate for the sensitivity of the detection method. Cells suitable for the present invention may be bacterial, yeast, or eukaryotic.

The assay methods to determine compound modulation of functional protease D-G can be in conventional laboratory format or adapted for high throughput. The term "high throughput" refers to an assay design that allows easy analysis of multiple samples simultaneously, and capacity for robotic manipulation. Another desired feature of high throughput assays is an assay

ORT1273

degree of cellular fluorescence. In flow cytometry, the cells are analyzed in a solution, whereas in cellular imaging techniques, a field of cells is compared for relative fluorescence.

A preferred detection means for secreted proteins that are enzymes such as alkaline phosphatase or proteases, would be fluorescent or colorimetric enzymatic assays. Fluorescent/luminescent/color substrates for alkaline phosphatase are commercially available and such assays are easily adaptable to high throughput multiwell plate screen format. Fluorescent energy transfer based assays are used for protease assays. Fluorophore and quencher molecules are incorporated into the two ends of the peptide substrate of the protease. Upon cleavage of the specific substrate, separation of the fluorophore and quencher allows the fluorescence to be detectable. When the secreted protein could be measure by radioactive methods, scintillation proximity technology could be used. The substrate of the protein of interest is immobilized either by coating or incorporation on a solid support that contains a fluorescent material. A radioactive molecule, brought in close proximity to the solid phase by enzyme reaction, causes the fluorescent material to become excited and emit visible light. Emission of visible light forms the basis of detection of successful ligand/target interaction, and is measured by an appropriate monitoring device. An example of a scintillation proximity assay is disclosed in United States Patent No. 4,568,649, issued February 4, 1986. Materials for these types of assays are commercially available from Dupont NEN® (Boston, Massachusetts) under the trade name FlashPlate™.

A preferred detection means where the endogenous gene results in phenotypic cellular structural changes is statistical image analysis the cellular morphology or intracellular phenotypic changes. For example, but not by way of limitation, and cell may change morphology such a rounding versus remaining flat against a surface, or may become growth-surface independent and thus resemble transformed cell phenotype well known in the art of tumor cell biology, or a cell may produce new outgrowths. Phenotypic changes that may occur intracellularly

ORT1273

include cytoskeletal changes, alteration in the entoplasmic reticulum/Golgi complex in response to new gene transcription, or production of new vesicles.

Where the endogenous gene encodes a soluble intracellular protein, changes in the endogenous gene may be measured by changes of the specific protein contained within the cell lysate. The soluble protein may be measured by the methods described herein.

The present invention is also directed to methods for screening for compounds that modulate the expression of DNA or RNA encoding protease D-G as well as the function of protease D-G protein *in vivo*. Compounds may modulate by increasing or attenuating the expression of DNA or RNA encoding protease D-G, or the function of protease D-G protein. Compounds that modulate the expression of DNA or RNA encoding protease D-G or the function of protease D-G protein may be detected by a variety of assays. The assay may be a simple "yes/no" assay to determine whether there is a change in expression or function. The assay may be made quantitative by comparing the expression or function of a test sample with the levels of expression or function in a standard sample. Modulators identified in this process are useful as therapeutic agents, and protease D-G.

Purification of protease D-G protein

Following expression of protease D-G in a recombinant host cell, protease D-G protein may be recovered to provide purified protease D-G in active form. Several protease D-G purification procedures are available and suitable for use (add references for purification of similar proteins that could be the basis of a purification scheme). As described above for purification of protease D-G from natural sources, recombinant protease D-G may be purified from cell lysates and extracts, or from conditioned culture medium, by various combinations of, or individual application of salt fractionation, ion exchange chromatography, size exclusion chromatography, hydroxylapatite adsorption chromatography and

ORT1273

hydrophobic interaction chromatography, lectin chromatography, and antibody/ligand affinity chromatography.

Recombinant protease D-G can be separated from other cellular proteins by use of an immunoaffinity column made with monoclonal or polyclonal antibodies specific for full length nascent protease D-G, polypeptide fragments of protease D-G or protease D-G subunits. The affinity resin is then equilibrated in a suitable buffer, for example phosphate buffered saline (pH 7.3), and the cell culture supernatants or cell extracts containing protease D-G or protease D-G subunits are slowly passed through the column. The column is then washed with the buffer until the optical density (A_{280}) falls to background, then the protein is eluted by changing the buffer condition, such as by lowering the pH using a buffer such as 0.23 M glycine-HCl (pH 2.6). The purified protease D-G protein is then dialyzed against a suitable buffer such as phosphate buffered saline.

Protein based assay

The present invention provides an *in vitro* protein assay method to detect compound modulation of protease D-G protein activity. The method comprises the steps;

- 1) contacting a compound, and function protease D-G protein, and
- 2) measuring a change to protease D-G function by the compound.

The amount of time necessary for cellular contact with the compound is empirically determined, for example, by running a time course with a known protease D-G modulator and measuring changes as a function of time.

this section is directed to protease assays

Methods for detecting compounds that modulate protease D-G proteolytic activity comprise combining a punitive modulating compound, functional protease D-G protein, and a suitable labeled substrate and monitoring an effect of the compound on the protease by changes in the amount of substrate either as a function of time or after a predefined period of time. Labeled substrates include, but are not limited to; substrate that is radiolabeled (Coolican et al. (1986). *J. Biol. Chem.* 261:4170-6),

fluorometric (Lonergan et al. (1995). *J. Food Sci.* 60:72-3, 78; Twining (1984). *Anal. Biochem.* 143:30-4) or colorimetric (Buroker-Kilgore and Wang (1993). *Anal. Biochem.* 208:387-92). Radioisotopes useful for use in the present invention include those well known in the art, specifically ^{125}I , ^{131}I , ^3H , ^{14}C , ^{35}S , ^{32}P , and ^{33}P .

- A preferred assay format useful for the method of the present invention is a FRET based method using peptide substrates that contain a fluorescent donor with either a quencher or acceptor that are separated by a peptide sequence encoding the protease D-G cleavage site. A fluorescent donor is a fluorogenic compound that can adsorb energy and transfers a portion of the energy to another compound. Examples of fluorescent donors suitable for use in the present invention include, but are not limited to, coumarins, xanthene dyes such as fluoresceins, rhodols, and rhodamines, resorufins, cyanine dyes bimanes, acridines, isoindols, dansyl dyes, aminophthalic hydrazides such as luminol and isoluminol derivatices, aminophthalimides, aminonaphthalimides, aminobenzofurans, aminoquinolines, dicanohydroquinones, and europium and terbium complexes and related compounds. A quencher is a compound that reduces the emission from the fluorescent donor when it is appropriately proximally located to the donor, and do not generally re-emit the energy in the form of fluorescence. Examples of such moieties include indigos, bezoquinones,

A preferred assay format useful for the method of the present invention is a FRET based method using peptide substrates that contain a fluorescent donor with either a quencher or acceptor that are separated by a peptide sequence encoding the protease D-G cleavage site. A fluorescent donor is a fluorogenic compound that can adsorb energy and transfers a portion of the energy to another compound. Examples of fluorescent donors suitable for use in the present invention include, but are not limited to, coumarins, xanthene dyes such as fluoresceins, rhodols, and rhodamines, resorufins, cyanine dyes bimanes, acridines, isoindols, dansyl dyes, aminophthalic hydrazides such as luminol and isoluminol derivatives, aminophthalimides, aminonaphthalimides, aminobenzofurans, aminoquinolines, dicanohydroquinones, and europium and terbium complexes and related compounds. A quencher is a compound that reduces the emission from the fluorescent donor when it is appropriately proximally located to the donor, and do not generally re-emit the energy in the form of fluorescence. Examples of such moieties include indigos, bezoquinones,

anthraquinones, azo compounds, nitro compounds, indoanilines, and di- and triphenylmethanes. A FRET method using a donor/quencher pair measures increased emission from the fluorescent donor as a function of protease D-G enzymatic activity upon the peptide substrate. Therefore a test compound th

20

Monospecific antibodies to protease D-G are purified from mammalian antisera containing antibodies reactive against protease D-G or are prepared as monoclonal antibodies reactive with protease D-G using the technique originally described by Kohler and Milstein, *Nature* 256: 495-497 (1975). Immunological techniques are well known in the art and described in, for example, Antibodies: A laboratory manual published by Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, ISBN 0879693142. Monospecific antibody as used herein is defined as a single antibody species or multiple antibody species with homogenous binding characteristics for protease D-G. Homogenous binding as

ORT1273

used herein refers to the ability of the antibody species to bind to a specific antigen or epitope, such as those associated with the protease D-G, as described above. protease D-G specific antibodies are raised by immunizing animals such as mice, rats, guinea pigs, rabbits, goats, horses and the like, with rabbits being
5 preferred, with an appropriate concentration of protease D-G either with or without an immune adjuvant.

Preimmune serum is collected prior to the first immunization. Each animal receives between about 0.001 mg and about 1000 mg of protease D-G associated with an acceptable immune adjuvant. Such acceptable adjuvants
10 include, but are not limited to, Freund's complete, Freund's incomplete, alum-precipitate, water in oil emulsion containing Corynebacterium parvum and tRNA. The initial immunization consists of protease D-G in, preferably, Freund's complete adjuvant at multiple sites either subcutaneously (SC), intraperitoneally (IP) or both. Each animal is bled at regular intervals, preferably weekly, to
15 determine antibody titer. The animals may or may not receive booster injections following the initial immunization. Those animals receiving booster injections are generally given an equal amount of the antigen in Freund's incomplete adjuvant by the same route. Booster injections are given at about three-week intervals until maximal titers are obtained. At about 7 days after each booster
20 immunization or about weekly after a single immunization, the animals are bled, the serum collected, and aliquots are stored at about -20°C.

Monoclonal antibodies (mAb) reactive with protease D-G are prepared by immunizing inbred mice, preferably Balb/c, with protease D-G. The mice are immunized by the IP or SC route with about 0.001 mg to about 1.0 mg, preferably
25 about 0.1 mg, of protease D-G in about 0.1 ml buffer or saline incorporated in an equal volume of an acceptable adjuvant, as discussed above. Freund's adjuvant is preferred, with Freund's complete adjuvant being used for the initial immunization and Freund's incomplete adjuvant used thereafter. The mice receive an initial immunization on day 0 and are rested for about 2 to about 30
30 weeks. Immunized mice are given one or more booster immunizations of about

ORT1273

0.001 to about 1.0 mg of protease D-G in a buffer solution such as phosphate buffered saline by the intravenous (IV) route. Lymphocytes, from antibody positive mice, preferably splenic lymphocytes, are obtained by removing spleens from immunized mice by standard procedures known in the art. Hybridoma cells are produced by mixing the splenic lymphocytes with an appropriate fusion partner, preferably myeloma cells, under conditions that will allow the formation of stable hybridomas. Fusion partners may include, but are not limited to: mouse myelomas P3/NS1/Ag 4-1; MPC-11; S-194 and Sp2/0, with Sp2/0 being generally preferred. The antibody producing cells and myeloma cells are fused in polyethylene glycol, about 1000 mol. wt., at concentrations from about 30% to about 50%. Fused hybridoma cells are selected by growth in hypoxanthine, thymidine and aminopterin supplemented Dulbecco's Modified Eagles Medium (DMEM) by procedures known in the art. Supernatant fluids are collected from growth positive wells on about days 14, 18, and 21 and are screened for antibody production by an immunoassay such as solid phase immunoradioassay (SPIRA) using protease D-G as the antigen. The culture fluids are also tested in the Ouchterlony precipitation assay to determine the isotype of the mAb. Hybridoma cells from antibody positive wells are cloned by a technique such as the soft agar technique of MacPherson, Soft Agar Techniques, in Tissue Culture Methods and Applications, Kruse and Paterson, Eds., Academic Press, 1973 or by the technique of limited dilution.

Monoclonal antibodies are produced *in vivo* by injection of pristane primed Balb/c mice, approximately 0.5 ml per mouse, with about 1×10^6 to about 6×10^6 hybridoma cells at least about 4 days after priming. Ascites fluid is collected at approximately 8-12 days after cell transfer and the monoclonal antibodies are purified by techniques known in the art.

In vitro production of anti-protease D-G mAb is carried out by growing the hybridoma in tissue culture media well known in the art. High density *in vitro* cell culture may be conducted to produce large quantities of anti-protease D-G mAbs using hollow fiber culture techniques, air lift reactors, roller bottle, or

ORT1273

spinner flasks culture techniques well known in the art. The mAb are purified by techniques known in the art.

Antibody titers of ascites or hybridoma culture fluids are determined by various serological or immunological assays which include, but are not limited to, precipitation, passive agglutination, enzyme-linked immunosorbent antibody (ELISA) technique and radioimmunoassay (RIA) techniques. Similar assays are used to detect the presence of protease D-G in body fluids or tissue and cell extracts.

It is readily apparent to those skilled in the art that the above described methods for producing monospecific antibodies may be utilized to produce antibodies specific for protease D-G polypeptide fragments, or full-length nascent protease D-G polypeptide, or the individual protease D-G subunits. Specifically, it is readily apparent to those skilled in the art that monospecific antibodies may be generated which are specific for only one protease D-G subunit or the fully functional protease D-G protein. It is also apparent to those skilled in the art that monospecific antibodies may be generated that inhibit normal function of protease D-G protein.

Protease D-G antibody affinity columns are made by adding the antibodies to a gel support such that the antibodies form covalent linkages with the gel bead support. Preferred covalent linkages are made through amine, aldehyde, or sulfhydryl residues contained on the antibody. Methods to generate aldehydes or free sulfhydryl groups on antibodies are well known in the art; amine groups are reactive with, for example, N-hydroxysuccinimide esters.

The aberrant expression or regulation of proteolytic activity can result in numerous pathophysiological states. For example several bleeding disorders, resulting from genetic lesions, are known to be caused by the deficiency in any one of a number of active serine protease coagulation factors. Many cancerous cells and tumors over-express proteases, several of which have been identified as serine proteases. These enzymes are thought to facilitate tumor growth and/or

5 extracellular matrix degradation or by the specific activation of pro-hormone precursors into active growth regulators or chemoattractants. Thus it is easy to imagine how modulators of serine protease activity could have profound effects of various pathophysiological conditions.

10 *Kit compositions containing protease D-G specific reagents*

Kits containing protease D-G DNA or RNA, antibodies to protease D-G, or protease D-G protein may be prepared. Such kits are used to detect DNA which hybridizes to protease D-G DNA or to detect the presence of protease D-G protein or peptide fragments in a sample. Such characterization is useful for a variety of purposes including but not limited to forensic analyses, diagnostic applications, and epidemiological studies.

The DNA molecules, RNA molecules, recombinant protein and antibodies of the present invention may be used to screen and measure levels of protease D-G DNA, protease D-G RNA or protease D-G protein. The recombinant proteins, DNA molecules, RNA molecules and antibodies lend themselves to the formulation of kits suitable for the detection and typing of protease D-G. Such a kit would comprise a compartmentalized carrier suitable to hold in close confinement at least one container. The carrier would further comprise reagents such as recombinant protease D-G protein or anti-protease D-G antibodies suitable for detecting protease D-G. The carrier may also contain a means for detection such as labeled antigen or enzyme substrates or the like.

Gene therapy

ORT1273

Nucleotide sequences that are complementary to the protease D-G encoding DNA sequence can be synthesized for antisense therapy. These antisense molecules may be DNA, stable derivatives of DNA such as phosphorothioates or methylphosphonates, RNA, stable derivatives of RNA such as 2'-O-alkylRNA, or other protease D-G antisense oligonucleotide mimetics. protease D-G antisense molecules may be introduced into cells by microinjection, liposome encapsulation or by expression from vectors harboring the antisense sequence. protease D-G antisense therapy may be particularly useful for the treatment of diseases where it is beneficial to reduce protease D-G activity.

protease D-G gene therapy may be used to introduce protease D-G into the cells of target organisms. The protease D-G gene can be ligated into viral vectors that mediate transfer of the protease D-G DNA by infection of recipient host cells. Suitable viral vectors include retrovirus, adenovirus, adeno-associated virus, herpes virus, vaccinia virus, polio virus and the like. Alternatively, protease D-G DNA can be transferred into cells for gene therapy by non-viral techniques including receptor-mediated targeted DNA transfer using ligand-DNA conjugates or adenovirus-ligand-DNA conjugates, lipofection membrane fusion or direct microinjection. These procedures and variations thereof are suitable for *ex vivo* as well as *in vivo* protease D-G gene therapy. protease D-G gene therapy may be particularly useful for the treatment of diseases where it is beneficial to elevate protease D-G activity. Protocols for molecular methodology of gene therapy suitable for use with the protease D-G gene is described in Gene Therapy Protocols, edited by Paul D. Robbins, Human press, Totawa NJ, 1996.

Pharmaceutical compositions

Pharmaceutically useful compositions comprising protease D-G DNA, protease D-G RNA, or protease D-G protein, or modulators of protease D-G receptor activity, may be formulated according to known methods such as by the admixture of a pharmaceutically acceptable carrier. Examples of such carriers and methods of formulation may be found in Remington's Pharmaceutical

ORT1273

Sciences. To form a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain an effective amount of the protein, DNA, RNA, or modulator.

Therapeutic or diagnostic compositions of the invention are administered to an individual in amounts sufficient to treat or diagnose disorders in which modulation of protease D-G-related activity is indicated. The effective amount may vary according to a variety of factors such as the individual's condition, weight, sex and age. Other factors include the mode of administration. The pharmaceutical compositions may be provided to the individual by a variety of routes such as subcutaneous, topical, oral and intramuscular.

The term "chemical derivative" describes a molecule that contains additional chemical moieties that are not normally a part of the base molecule. Such moieties may improve the solubility, half-life, absorption, etc. of the base molecule. Alternatively the moieties may attenuate undesirable side effects of the base molecule or decrease the toxicity of the base molecule. Examples of such moieties are described in a variety of texts, such as Remington's Pharmaceutical Sciences.

Compounds identified according to the methods disclosed herein may be used alone at appropriate dosages defined by routine testing in order to obtain optimal inhibition of the protease D-G receptor or its activity while minimizing any potential toxicity. In addition, co-administration or sequential administration of other agents may be desirable.

The present invention also has the objective of providing suitable topical, oral, systemic and parenteral pharmaceutical formulations for use in the novel methods of treatment of the present invention. The compositions containing compounds or modulators identified according to this invention as the active ingredient for use in the modulation of protease D-G can be administered in a wide variety of therapeutic dosage forms in conventional vehicles for administration. For example, the compounds or modulators can be administered in such oral dosage forms as tablets, capsules (each including timed release and

ORT1273

sustained release formulations), pills, powders, granules, elixirs, tinctures, solutions, suspensions, syrups and emulsions, or by injection. Likewise, they may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed as a protease D-G modulating agent.

The daily dosage of the products may be varied over a wide range from 0.01 to 1,000 mg per patient, per day. For oral administration, the compositions are preferably provided in the form of scored or unscored tablets containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, and 50.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.0001 mg/kg to about 100 mg/kg of body weight per day. The range is more particularly from about 0.001 mg/kg to 10 mg/kg of body weight per day. The dosages of the protease D-G receptor modulators are adjusted when combined to achieve desired effects. On the other hand, dosages of these various agents may be independently optimized and combined to achieve a synergistic result wherein the pathology is reduced more than it would be if either agent were used alone.

Advantageously, compounds or modulators of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds or modulators for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

ORT1273

For combination treatment with more than one active agent, where the active agents are in separate dosage formulations, the active agents can be administered concurrently, or they each can be administered at separately staggered times.

- 5 The dosage regimen utilizing the compounds or modulators of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound thereof employed. A physician or
10 veterinarian of ordinary skill can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentrations of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the
15 distribution, equilibrium, and elimination of a drug.

- In the methods of the present invention, the compounds or modulators herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected
20 with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

- For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically
25 acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or
30 sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like.

Lubricants used in these dosage forms include, without limitation, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

Topical preparations containing the active drug component can be admixed with a variety of carrier materials well known in the art, such as, e.g., alcohols, aloe vera gel, allantoin, glycerine, vitamin A and E oils, mineral oil, PPG2 myristyl propionate, and the like, to form, e.g., alcoholic solutions, topical cleansers, cleansing 15 creams, skin gels, skin lotions, and shampoos in cream or gel formulations.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds or modulators of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinyl-pyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxy-ethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds or modulators of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for

ORT1273

example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydro-pyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

For oral administration, the compounds or modulators may be administered in capsule, tablet, or bolus form or alternatively they can be mixed in the animals feed. The capsules, tablets, and boluses are comprised of the active ingredient in combination with an appropriate carrier vehicle such as starch, talc, magnesium stearate, or di-calcium phosphate. These unit dosage forms are prepared by intimately mixing the active ingredient with suitable finely-powdered inert ingredients including diluents, fillers, disintegrating agents, and/or binders such that a uniform mixture is obtained. An inert ingredient is one that will not react with the compounds or modulators and which is non-toxic to the animal being treated. Suitable inert ingredients include starch, lactose, talc, magnesium stearate, vegetable gums and oils, and the like. These formulations may contain a widely variable amount of the active and inactive ingredients depending on numerous factors such as the size and type of the animal species to be treated and the type and severity of the infection. The active ingredient may also be administered as an additive to the feed by simply mixing the compound with the feedstuff or by applying the compound to the surface of the feed. Alternatively the active ingredient may be mixed with an inert carrier and the resulting composition may then either be mixed with the feed or fed directly to the animal. Suitable inert carriers include corn meal, citrus meal, fermentation residues, soya grits, dried grains and the like. The active ingredients are intimately mixed with these inert carriers by grinding, stirring, milling, or tumbling such that the final composition contains from 0.001 to 5% by weight of the active ingredient.

The compounds or modulators may alternatively be administered parenterally via injection of a formulation consisting of the active ingredient dissolved in an inert liquid carrier. Injection may be either intramuscular,

ORT1273

intraruminal, intratracheal, or subcutaneous. The injectable formulation consists of the active ingredient mixed with an appropriate inert liquid carrier. Acceptable liquid carriers include the vegetable oils such as peanut oil, cotton seed oil, sesame oil and the like as well as organic solvents such as solketal, glycerol formal and the like. As an alternative, aqueous parenteral formulations may also be used. The vegetable oils are the preferred liquid carriers. The formulations are prepared by dissolving or suspending the active ingredient in the liquid carrier such that the final formulation contains from 0.005 to 10% by weight of the active ingredient.

Topical application of the compounds or modulators is possible through the use of a liquid drench or a shampoo containing the instant compounds or modulators as an aqueous solution or suspension. These formulations generally contain a suspending agent such as bentonite and normally will also contain an antifoaming agent. Formulations containing from 0.005 to 10% by weight of the active ingredient are acceptable. Preferred formulations are those containing from 0.01 to 5% by weight of the instant compounds or modulators.

The following examples illustrate the present invention without, however, limiting the same thereto.

EXAMPLE 1

Plasmid Manipulations

All molecular biological methods were in accordance with those previously described (Maniatis et al. (1989). 1-1626). Oligonucleotides were purchased from Ransom Hill Biosciences (Ransom Hill, CA) and all restriction endonucleases and other DNA modifying enzymes were from New England Biolabs (Beverly, MA) unless otherwise specified. The protease D-G expression construct was made in the baculovirus expression vector pFastBac1 (Life Technologies, Gaithersburg, MD) as described below. All construct

manipulations were confirmed by dye terminator cycle sequencing using Allied Biosystems 377 fluorescent sequencers (Perkin Elmer, Foster City, CA).

A recombinant phage containing the protease D-G cDNA was isolated from a human small intestine library (Clontech, Palo Alto, CA). The insert was subjected to sequence analysis and it was found to contain an open reading frame of 1305 nucleotides excluding the TAA stop codon (SEQ ID NO.:1), which had homology to S1 serine proteases. Significantly, the open reading frame is likely to be authentic since it is preceded by an in-frame TGA stop codon at position 157. This clone is also likely to contain the entire 3' untranslated since a putative polyadenylation sequence (ATTAAA) with a good match to the known sequence was (AATAAA) was also identified just upstream of a poly A stretch. The deduced open reading frame encodes a preproD-G protein of 435 amino acids (SEQ ID NO.:2), with an estimated molecular mass (M_r) of about 48-Kd, and a strong homology to other serine proteases. Additional sequence analysis of the protease D-G amino acid sequence predicted a transmembrane segment near the amino terminus (residues 31-52 in SEQ ID NO.:2), suggesting that this novel cDNA encoded a type II transmembrane serine protease. Homology searches of the Genbank database with the protease D-G cDNA indicated that this was a novel cDNA had closest similarity to the cloned serine proteases TMPRSS2 (Paoloni-Giacobino et al. (1997). *Genomics* 44:309-320) and hepsin (Leytus et al. (1988). *Biochemistry* 27:1067-74), which are also type II integral membrane proteases. The zymogen activation sequence is very similar to that of other S1 serine proteases and predicts a mature protein of 233 amino acids. The catalytic triad residues H, D and S of protease D-G are located at positions 243, 339 and 385, respectively (using the methionine initiator of the prepro D-G sequence as number one). A phylogenetic tree of the deduced protease D-G amino acid sequence with other members of the S1 serine protease family was generated by the Clustal W program (Higgins and Sharp (1989). *Comput. Appl. Biosci.* 5:151-3) shown in Figure 2 as determined using the MegAlign 3.1.7 program (DNASTAR Inc., Madison, WI).

5 A recombinant phage containing the protease D-G cDNA was isolated from a human small intestine library (Clontech, Palo Alto, CA). The insert was subjected to sequence analysis and it was found to contain an open reading frame of 1305 nucleotides excluding the TAA stop codon (SEQ ID NO.:1), which had homology to S1 serine proteases. Significantly, the open reading frame is likely to be authentic since it is preceded by an in-frame TGA stop codon at position 157. This clone is also likely to contain the entire 3' untranslated since a putative polyadenylation sequence (ATTAAA) with a good match to the known sequence was (AATAAA) was also identified just upstream of a poly A stretch. The deduced open reading frame encodes a preproD-G protein of 435 amino acids (SEQ ID NO.:2), with an estimated molecular mass (M_r) of about 48-Kd, and a strong homology to other serine proteases. Additional sequence analysis of the protease D-G amino acid sequence predicted a transmembrane segment near the amino terminus (residues 31-52 in SEQ ID NO.:2), suggesting that this novel cDNA encoded a type II transmembrane serine protease. Homology searches of the Genbank database with the protease D-G cDNA indicated that this was a novel cDNA had closest similarity to the cloned serine proteases TMPRSS2 (Paoloni-Giacobino et al. (1997). *Genomics* 44:309-320) and hepsin (Leytus et al. (1988). *Biochemistry* 27:1067-74), which are also type II integral membrane proteases. The zymogen activation sequence is very similar to that of other S1 serine proteases and predicts a mature protein of 233 amino acids. The catalytic triad residues H, D and S of protease D-G are located at positions 243, 339 and 385, respectively (using the methionine initiator of the prepro D-G sequence as number one). A phylogenetic tree of the deduced protease D-G amino acid sequence with other members of the S1 serine protease family was generated by the Clustal W program (Higgins and Sharp (1989). *Comput. Appl. Biosci.* 5:151-3) shown in Figure 2 as determined using the MegAlign 3.1.7 program (DNASTAR Inc., Madison, WI).

ORT1273

EXAMPLE 2

Tissue Distribution of The Protease D-G mRNA

We employed a highly sensitive PCR profiling technique to identify the tissue distribution of protease D-G mRNA. For this application, several human cDNA libraries (all were from Clontech, (Palo Alto, CA) except the CHRF-288 megakaryocytic cell line and human gel filtered platelet libraries which we constructed using the ZAP Express cDNA system (Stratagene, La Jolla, CA). The PCR primers for the profiling analysis were as follows:

10 SEQ.ID.NO.3: 5'- ACAGCCTCAGCATTCTTGG -3'

SEQ.ID.NO.4: 5'- TCTTGCTCTAGTAGGCTTGG -3'

Briefly, the 50µl PCR reactions used 1 µl of diluted phage stock ($\sim 10^8$ to 10^{10} pfu/ml) from each of the cDNA libraries tested. Reactions were initially denatured at 94 °C for 5 min. and subjected to 35 cycles of 94 °C for 20 sec.; 56 °C for 20 sec.; and then 72 °C for 30 sec. followed by a final 72 °C elongation for 10 min. A nested primer probe of the sequence

SEQ.ID.NO.5: 5'-

TTGGTGGCTCCCAGCATCCCAGGGAGAGACACAGCCCACTG -3' was radiolabeled using gamma ³²P-ATP and T4 polynucleotide kinase (Life Technologies, Gaithersburg, MD) and unincorporated label was removed, following the reaction, using a QIAquick nucleotide removal column (Qiagen, Valencia, CA). The ³²P end-labeled nested primer probe (1X10⁵ cpm) was combined with 10 µl of each sample following the PCR reaction. The PCR product-probe mixtures were denatured at 94 °C for 5 min.; hybridized at 60 °C for 15 minutes, and cooled to 4 °C. The annealed samples (10 µl) were electrophoresed in 6% Tris-Borate-EDTA non-denaturing polyacrylamide gels (Novex), dried and exposed by autoradiography. A PCR profile of the cDNA libraries used in Figure 3 with beta-actin PCR primers and labeled nested primer probe produced a beta-actin PCR product in all samples examined.

As seen in Figure 3, the distribution of protease D-G mRNA is highly
30 restricted to specific tissues and cell types. The tissue types expressing the protease

D-G transcript are epidermis, fibroblasts, keratinocytes, colon, small intestine, stomach, lung, kidney, bone marrow, lymph node, thymus, ovary, prostate, uterus and spinal cord. Of particular significance is that D-G protease mRNA is not expressed in pancreas or liver, tissues normally found to express numerous serine protease genes.

Construct Generation For The Expression of Active Protease D-G

The purified plasmid DNA of the full length protease D-G cDNA was used as a template in a 100 µl preparative PCR reaction using the Native Pfu Polymerase (Stratagene, La Jolla, CA) in accordance with the manufacturer's recommendations. The primers used

SEQ.ID.NO.6: D-G Xba-U 5'- ATGCTCTAGATGTGGATTCTTGGCCTTGGC
-3' SEQ.ID.NO.7: D-G Xba-L 5'-

ORT1273

GATGTCTAGACAGCTCAGCCTTCCAGACATTG -3' contained Xba I cleavable ends, and were designed to flank the catalytic domain of protease D-G and generate the protease D-G Xba I catalytic cassette. The preparative PCR reaction was run at 18 cycles of 94°C for 30 sec.; 60°C for 30 sec; 72°C for 2.0 min.

The preparative PCR product was phenol/CHCl₃ (1:1) extracted once, CHCl₃ extracted, and then EtOH precipitated with glycogen (Boehringer Mannheim Corp., Indianapolis, IN) and carrier. The precipitated pellet was rinsed with 70 % EtOH, dried by vacuum, and resuspended in 80 ul H₂O, 10 ul 10 restriction buffer number 2 and 1 ul 100x BSA (New England Biolabs, Beverly, MA). The product was digested for 3 hr. at 37 °C with 200 units Xba I restriction enzyme (New England Biolabs, Beverly, MA). The Xba I digested product was phenol/CHCl₃ (1:1) extracted once, CHCl₃ extracted, EtOH precipitated, rinsed with 70 % EtOH, and dried by vacuum. For purification from contaminating template plasmid DNA, the product was electrophoresed through 1.0 % low melting temperature agarose (Life Technologies, Gaithersburg, MD) gels in TAE buffer (40 mM Tris-Acetate, 1 mM EDTA pH 8.3) and excised from the gel. An aliquot of the excised product was then used for in-gel ligations with the Xba I digested, dephosphorylated and gel purified, zymogen activation vector described above. Clones containing the D-G Xba cassette, inserted in the correct orientation to generate the construct PFEK-protease D-G-6XHIS-TAG 64, were confirmed by sequence analyses to ensure that the proper translational register with respect to the NH₂-terminal PFEK prepro sequence and C-terminal 6XHIS affinity tag was maintained.

25

EXAMPLE 4

Expression of Recombinant Protease D-G

The recombinant bacmid containing the PFEK-protease D-G-6XHIS construct was prepared from bacterial transformation, selection, growth, purification and PCR confirmation in accordance with the manufacturer's recommendations.

30

ORT1273

Cultured Sf9 insect cells (ATCC CRL-1711) were transfected with purified bacmid DNA and several days later, conditioned media containing recombinant PFEK- protease D-G-6XHIS baculovirus was collected for viral stock amplification. Sf9 cells growing in Sf-900 II SFM at a density of 2×10^6 /ml were
5 infected at a multiplicity of infection of 2 at 27 °C for 80 hours, and media was collected and concentrated for purification of PFEK- protease D-G-6XHIS.

EXAMPLE 5

Purification, and Activation of Recombinant Protease D-G

Culture supernatants from baculovirus infected Sf9 cells expressing PFEK-
10 D-G-6XHIS were concentrated and desalted at 4 °C using a Centricon Plus-80 Biomax-8 concentrator (Millipore, Marlborough, MA). Ni-NTA (150 µl of a 50 % slurry/per 100 µg of zymogen) (Qiagen, Valencia, CA) was added to 5 ml the concentrated sample and mixed by shaking at 4 °C for 60 min. The zymogen-bound resin was washed 3 times with wash buffer [10 mM Tris-HCl (pH 8.0), 300
15 mM NaCl, and 15 mM imidazole], followed by with a 1.5 ml wash with ds H₂O. Zymogen cleavage was carried out by adding enterokinase (10 U per 50 µg of zymogen) (Novagen, Inc., Madison WI; or Sigma, St. Louis, MO) to the zymogen-bound Ni-NTA beads in a small volume at room temperature overnight with gentle shaking in a buffer containing 20 mM Tris-HCl (pH 7.4), 50 mM
20 NaCl, and 2.0 mM CaCl₂. The resin was then washed twice with 1.5 ml wash buffer. The activated protease D-G-6XHIS was eluted with elution buffer [20 mM Tris-HCl (pH 7.8), 250 mM NaCl, and 250 mM imidazole]. Eluted protein concentration was determined by a Micro BCA Kit (Pierce, Rockford, IL) using bovine serum albumin as a standard.

25

Electrophoresis and Western Blotting Detection of Recombinant Proteases D-G

Samples of the purified PFEK-protease D-G-6XHIS zymogen or activated protease D-G-6XHIS, denatured in the presence of the reducing agent dithiothreitol (DTT), were analyzed by SDS-PAGE (Bio Rad, Hercules CA) stained with
30 Coomassie Brilliant Blue. For Western blotting, gels were electrotransferred to

0060745:063000

ORT1273

Hybond ECL membranes (Amersham, Arlington Heights, IL). The FLAG-tagged PFEK-protease D-G-6XHIS zymogen expressed from infected Sf9 cells was detected with anti-Flag M2 antibody (Babco, Richmond, CA). The secondary antibody was a goat-anti-mouse IgG (H+L), horseradish peroxidase-linked F(ab')₂ fragment, (Boehringer Mannheim Corp., Indianapolis, IN) and was detected by the ECL kit (Amersham, Arlington Heights, IL).

EXAMPLE 6

Chromogenic Assay of Activated Recombinant Proteases D-G

Amidolytic activities of the activated serine proteases are monitored by release of para-nitroaniline (pNA) from synthetic substrates that are commercially available (Bachem California Inc., Torrance, PA; American Diagnostica Inc., Greenwich, CT; Kabi Pharmacia Hepar Inc., Franklin, OH). Assay mixtures contain chromogenic substrates in 500 μ M and 10 mM TRIS-HCl (pH 7.8), 25 mM NaCl, and 25 mM imidazole. Release of pNA is measured over 120 min at 37 °C on a micro-plate reader (Molecular Devices, Menlo Park, CA) with a 405 nm absorbance filter. The initial reaction rates (V_{max} , mOD/min) are determined from plots of absorbance versus time using Softmax (Molecular Devices, Menlo Park, CA). The specific activities (nmole pNA produced /min/ μ g protein) of the activated protease D-G-6XHIS for the various substrates are presented in Table 1. No measurable chromogenic amidolytic activity was detected with the purified unactivated PFEK-protease D-G-6XHIS zymogen.

TABLE 1
SPECIFIC ACTIVITY TABLE

ORT1273

Chromogenic Substrates	Specific Activity
H-D-Pro-HHT-Arg-pNA	0.046±0.001
H-D-Lys(CBO)-Pro-Arg-pNA	0.076±0.008
Z-Phe-Arg-pNA	0.116±0.006
H-D-Val-Leu-Arg-pNA	0.025±0.003
H-D-Val-Leu-Lys-pNA	0.034±0.003
Suc-Ala-Ala-Pro-Phe-pNA	N.A.
Meo-Suc-Ala-Ala-Pro-Val-pNA	N.A.
	N.A. = No Activity

Table 1 - The specific activity (nmole pNA produced /min/ug protein) of recombinant activated protease D-G-6XHIS, determined for the various substrates analyzed, is shown.

5

Compounds that modulate a serine protease of the present invention are identified through screening for the acceleration, or more commonly, the inhibition of the proteolytic activity. Although in the present case chromogenic activity is monitored by an increase in absorbance, fluorogenic assays or other methods such as FRET to measure proteolytic activity as mentioned above, can be employed. Compounds are dissolved in an appropriate solvent, such as DMF, DMSO, methanol, and diluted in water to a range of concentrations usually not exceeding 100 μ M and are typically tested, though not limited to, a concentration of 1000-fold the concentration of protease. The compounds are then mixed with the protein stock solution, prior to addition to the reaction mixture. Alternatively, the protein and compound solutions may be added independently to the reaction mixture, with the compound being added either prior to, or immediately after, the addition of the protease D-G protein.

20

During the course of these investigations a submission in Genbank was identified (Genbank accession number AF216312) which is similar but distinct from

the sequence described herein. Although the exact significance of the discrepancy between the protease D-G cDNA and the AF216312 sequence is not fully understood at this time, it may be a result of alternative splicing near the initiator ATG, thereby generating distinct coding sequences and consequently distinct proteins. Below is a

5 GAP alignment between the nucleic acid sequences of the protease D-G cDNA
(SEQ.ID.NO.:1) described herein on top and in upper case, and the AF216312
sequence indicated below and in lower case.

The Genbank explanatory information is reproduced as follows:

LOCUS AF216312 2079 bp mRNA

10 DEFINITION Homo sapiens type II membrane serine protease mRNA, complete cds.

ACCESSION AF216312 VERSION AF216312.1 GI:6911218

SOURCE human.

ORGANISM Homo sapiens Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

15 REFERENCE 1 (bases 1 to 2079)

AUTHORS Smeekens, S.S., Lorimer, D.D., Wang, E., Hou, J. and Linnevers, C.

TITLE MT-SP2, a novel type II membrane serine protease expressed in trachea, colon, and small intestine: identification, cloning, and chromosomal localization

JOURNAL Unpublished

20 REFERENCE 2 (bases 1 to 2079)

AUTHORS Smeekens, S.S., Lorimer, D.D., Wang, E., Hou, J. and Linnevers, C.

TITLE Direct Submission

JOURNAL Submitted (14-DEC-1999) Axys Pharmaceuticals, Inc, 180 Kimball Way,
South San Francisco, CA 94080, USA

25

SEQ.ID.NO.:1 x AF216312.Seq

51 CACTCCTGGAATACACAGAGAGAGGCAGCTTGCTCAGCGGACAAGGA 100
 1gagaggcagcagcttgttcagcggacaagga 31
 101 TGCTGGGCGTGAGGGACCAAGGCCTGCCCTGCACTCGGGCCTCCTCCAGC 150

	32	tgctggggcgtgagggaccaaggcctgccctgcactcgggcctcctccagc	81
	151	CAGTGCTGACCAGGGACTTCTGACCTGCTGGCCAGCCAGGACCTGTGTGG	200
5	82	cagtgtgaccagggacttctgacctgctggccagccaggacctgtgtgg	131
	201	GGAGGCCCTCCTGCTGCCTTGGGGTGACAATCTCAGCTCCAGGCTACAGG	250
10	132	ggaggccctcctgctgccttgggggtgacaatctcagctccaggctacagg	181
	251	GAGACCGGGAGGATCACAGAGCCAGCAT.....GGATCCTGACAGTGAT	294
	182	gagaccgggaggatcacagagccagcatggtacaggatcctgacagtgat	231
15	295	CAACCTCTGAACAGCCTCGATGTCAAACCCCTGCGCAAACCCGTATCCC	344
	232	caacctctgaacagcctcgatgtcaaaccctgcgcaaaccctgtatccc	281
20	345	CATGGAGACCTTCAGAAAG.GTGGGGATCCCCATCATCATAGCACTACTG	393
	282	catggagaccttcagaaagtgtggggatccccatcatcatagcactactg	331
	394	AGCCTGGCGAGTATCATCATTGTGGTTGTCTCATCAAGGTGATTCTGGA	443
25	332	agcctggcgagtatcatcattgtggttgtcctcatcaaggatgattctgga	381
	444	TAAATACTACTTCCTCTGCGGGCAGCCTCTCCACTTCATCCCGAGGAAGC	493
30	382	taaatactacttcctctgcgggcagcctctccacttcatcccaggaagc	431
	494	AGCTGTGTGACGGAGAGCTGGACTGTCCCTTGGGGGAGGACGAGGAGCAC	543
	432	agctgtgtgacggagagctggactgtcccttgggggaggacgaggagcac	481
35	544	TGTGTCAAGAGCTTCCCCGAAGGGCCTGCAGTGGCAGTCCGCCTCTCCAA	593
	482	tgtgtcaagagcttccccgaagggcctgcagtggcagtcgcgctctccaa	531
40	594	GGACCGATCCCACTGCAGGTGCTGGACTCGGCCACAGGGAAGTGGTTCT	643
	532	ggaccgatccacactgcaggtgctggactcggccacagggaaactggttct	581
	644	CTGCCTGTTTCGACAACTTCACAGAAGCTCTCGCTGAGACAGCCTGTAGG	693
45	582	ctgcctgtttcgacaacttcacagaagctctcgctgagacagcctgtagg	631
	694	CAGATGGGCTACAGCAGCAAACCCACTTTTCAGAGCTGTGGAGATTGGCCC	743
50	632	cagatgggctacagcagcaaaccactttcagagctgtggagattggccc	681

	744	AGACCAGGATCTGGATGTTGTTGAAATCACAGAAAACAGCCAGGAGCTTC	793
	682	agaccaggatctggatggtgttgaaatcacagaaaacagccaggagcttc	731
5	794	GCATGCGGAACTCAAGTGGGCCCTGTCTCTCAGGCTCCCTGGTCTCCCTG	843
	732	gcatgcggaactcaagtgggccctgtctctcaggctccctggtctccctg	781
10	844	CACTGTCTTGCCTGTGGGAAGAGCCTGAAGACCCCCCGTGTGGTGGGTGG	893
	782	cactgtcttgcctgtgggaagagcctgaagaccccccggtgtggtgggtgg	831
	894	GGAGGAGGCCCTCTGTGGATTCTTGGCCTTGGCAGGTCAGCATCCAGTACG	943
15	832	ggaggaggcctctgtggattcttggccttggcaggtcagcatccagtacg	881
	944	ACAAACAGCACGTCTGTGGAGGGAGCATCCTGGACCCCCACTGGGTCTCTC	993
	882	acaaacagcacgtctgtggagggagcatcctggacccccactgggtcctc	931
20	994	ACGGCAGCCCCTGCTTCAGGAAACATACCGATGTGTTCAACTGGAAGGT	1043
	932	acggcagcccactgcttcaggaaacataccgatgtgttcaactggaaggt	981
25	1044	GCGGGCAGGCTCAGACAAACTGGGCAGCTTCCCATCCCTGGCTGTGGCCA	1093
	982	gcgggcaggctcagacaaactgggcagcttcccatccctggctgtggcca	1031
30	1094	AGATCATCATCATTGAATTCAACCCCATGTACCCCAAAGACAATGACATC	1143
	1032	agatcatcatcattgaattcaaccccattgtaccccaaagacaatgacatc	1081
	1144	GCCCTCATGAAGCTGCAGTTCCTCACTTTCTCAGGCACAGTCAGGCC	1193
35	1082	gccctcatgaagctgcagttccctactcactttctcaggcacagttaggcc	1131
	1194	CATCTGTCTGCCCTTCTTTGATGAGGAGCTCACTCCAGCCACCCCACTCT	1243
	1132	catctgtctgcccttctttgatgaggagctcactccagccaccccactct	1181
40	1244	GGATCATTGGATGGGGCTTTACGAAGCAGAATGGAGGGAAGATGTCTGAC	1293
	1182	ggatcattggatggggctttacgaagcagaatggaggggaagatgtctgac	1231
45	1294	ATACTGCTGCAGGCGTCAGTCCAGGTCATTGACAGCACACGGTGCAATGC	1343
	1232	atactgctgcaggcgtcagtcagggtcattgacagcacacggtgcaatgc	1281
50	1344	AGACGATGCGTACCTGGGGGAAGTCACCGAGAAGATGATGTGTGCAGGCA	1393

	1282	agacgatgcgtaccaggggggaagtaccgcagaagatgatgtgtgcaggca	1331
	1394	TCCCGGAAGGGGGTGTGGACACCTGCCAGGGTGACAGTGGTGGGCCCTG	1443
5	1332	tcccggaaggggggtgtggacacctgccagggtagacagtgggtgggcccctg	1381
	1444	ATGTACCAATCTGACCAGTGGCATGTGGTGGGCATCGTTAGCTGGGGCTA	1493
10	1382	atgtaccaatctgaccagtggcatgtgggtgggcatcgttagctggggcta	1431
	1494	TGGCTGCGGGGGCCCCGAGCACCCCAGGGGTATACACCAAGGTCTCAGCCT	1543
	1432	tggctgcggggggcccgagcaccacaggagtatacaccaaggtctcagcct	1481
15	1544	ATCTCAACTGGATCTACAATGTCTGGAAGGCTGAGCTGTAATGCTGCTGC	1593
	1482	atctcaactggatctacaatgtctggaaggctgagctgtaatgctgctgc	1531
	1594	CCCTTTGCAGTGCTGGGAGCCGCTTCCTTCCTGCCCTGCCACCTGGGGA	1643
20	1532	ccctttgcagtgtctgggagccgcttccttcctgcctgcccacctgggga	1581
	1644	TCCCCCAAAGTCAGACACAGAGCAAGAGTCCCCTTGGGTACACCCCTCTG	1693
25	1582	tcccccaaagtcagacacagagcaagagtccttgggtacaccctctg	1631
	1694	CCCACAGCCTCAGCATTCTTGGAGCAGCAAAGGGCCTCAATTCTATAA	1743
30	1632	cccacagcctcagcatttcttggagcagcaaagggcctcaattcctgtaa	1681
	1744	GAGACCCTCGCAGCCCAGAGGCGCCAGAGGAAGTCAGCAGCCCTAGCTC	1793
	1682	gagaccctcgcagcccagaggcgccagaggaagtcagcagccctagctc	1731
35	1794	GGCCACACTTGGTGCTCCCAGCATCCCAGGGAGAGACACAGCCCACTGAA	1843
	1732	ggccacacttgggtgctcccagcatcccaggagagacacagcccaactgaa	1781
	1844	CAAGGTCTCAGGGGTATTGCTAAGCCAAGAAGGAACCTTCCCACACTACT	1893
40	1782	caaggtctcaggggtattgctaagccaagaaggaactttcccacactact	1831
	1894	GAATGGAAGCAGGCTGTCTTGTAAGCCAGCCAGATCACTGTGGGCTGGAGA	1943
45	1832	gaatggaagcaggctgtcttgtaaaagcccagatcactgtgggctggaga	1881
	1944	GGAGAAGGAAAGGGTCTGCGCCAGCCCTGTCCGTCTTCACCCATCCCCAA	1993
50	1882	ggagaaggaaagggctctgcgccagccctgtccgtcttcacccatccccaa	1931

```

1994 GCCTACTAGAGCAAGAAACCAGTTGTAATATAAAATGCACTG.CCTACTG 2042
      |||||||||||||||||||||||||||||||||||||||||||||
1932 gcctactagagcaagaaaccagttgtaatatataaaatgcactgccctactg 1981
      .
5 2043 TTGGTATGACTACCGTTACCTACTGTTGTCATTGTTATTACAGCTATGGC 2092
      |||||||||||||||||||||||||||||||||||||||||||||
      .
1982 ttggtatgactaccgttacctactgttgtcattgttattacagctatggc 2031
      .
      .
10 2093 CACTATTATTAAAGAGCTGTGTAACATCA..... 2121
      |||||||||||||||||||||||||||
2032 CACTATTATTAAAGAGCTGTGTAACATCAAAAAAAAAAAAAAAAAAAAA 2079

```

Below is a GAP alignment between the amino acid sequences of the protease D-G cDNA (SEQ.ID.NO.:2) described herein, on top with the predicted transmembrane domain in lower case, and that encoded by the AF216312 sequence indicated below.

20	1	MDPDS	DQPLNSLDVKPLRKPRIPMETFRKV	gipiiiallsiasiiivvvl	50
	1	MSNPCANPVSPWRPSESVGIPIIIALLSLASIIIVVVL		38
25	51	ikVILD	KYYFLCGQPLHFIPRKQLCDGELDCPLGEDEEHCVKSFPEGPAV		100
	39	IKVILD	KYYFLCGQPLHFIPRKQLCDGELDCPLGEDEEHCVKSFPEGPAV		88
	101	AVRLSKDRSTLQVLDSATGNWF	SACFDNFTEALAE	TACRQMGYSSKPTFR	150
30	89	AVRLSKDRSTLQVLDSATGNWF	SACFDNFTEALAE	TACRQMGYSSKPTFR	138
	151	AVEIGPDQDLDVVEITENSQEL	MRNSSGPCLSGSLVSLHCLACGKSLKT		200
	139	AVEIGPDQDLDVVEITENSQEL	MRNSSGPCLSGSLVSLHCLACGKSLKT		188
35					
	201	PRVVG	GEEASVDSWPWQVSIQYDKQHVC	GSILDPHWVLTAAHCFRKHTD	250
	189	PRVVG	GEEASVDSWPWQVSIQYDKQHVC	GSILDPHWVLTAAHCFRKHTD	238
40	251	VF	NWKVRAGSDKLG	SFPSLAVAKIIIIIEFNPMYPKDNDIALMKLQFPLTF	300
	239	VF	NWKVRAGSDKLG	SFPSLAVAKIIIIIEFNPMYPKDNDIALMKLQFPLTF	288
45	301	SG	TVRPICLPFFDEELTPATPLWII	IGWGFTKQNGGKMSDILLQASVQVID	350

```

289 SGTVRPICLPFFDEELTPATPLWIIIGWGFTKQNGGKMSDILLQASVQVID 338
      .           .           .           .
351 STRCNADDAYQGEVTEKMMCAGIPEGGVDTTCQGDSGGPLMYQSDQWHVVG 400
      |||||
5  339 STRCNADDAYQGEVTEKMMCAGIPEGGVDTTCQGDSGGPLMYQSDQWHVVG 388
      .           .           .
401 IVSWGYGCGGPSTPGVYTKVSAYLNWIYNVWKAEL 435
      |||||
10 389 IVSWGYGCGGPSTPGVYTKVSAYLNWIYNVWKAEL 423

```